

Location: Cattai Wetlands


Survey method: Fixed walk around main lagoon, duration 1.5-2 hours

N^o of species recorded: 143

Date this list prepared: 1 September 2018

N^o of surveys conducted (up to the preparation date): 62

Comment: The focus of the survey is waterbirds; any bush birds encountered are recorded but non-calling birds in woodlands mostly are over-looked



Alan Stuart

81 Queens Rd
New Lambton NSW 2305 Australia
Mobile: +61 409 978171
Email: alanstuart400@gmail.com
Web: www.thinkingaboutbirds.com

Species recorded, and their reporting rates

| | | | |
|---------------------------|-------|------------------------------|-------|
| Wandering Whistling-Duck | 8.1% | Australian Pelican | 22.6% |
| Pink-eared Duck | 3.2% | Nankeen Night-Heron | 1.6% |
| Black Swan | 80.7% | White-necked Heron | 16.1% |
| Hardhead | 53.2% | Great Egret | 30.7% |
| Australasian Shoveler | 22.6% | Intermediate Egret | 3.2% |
| Pacific Black Duck | 91.9% | White-faced Heron | 17.7% |
| Grey Teal | 69.4% | Australian White Ibis | 8.1% |
| Chestnut Teal | 61.3% | Straw-necked Ibis | 6.5% |
| Australian Wood Duck | 4.8% | Royal Spoonbill | 4.8% |
| Brown Quail | 9.7% | Little Pied Cormorant | 46.8% |
| King Quail | 1.6% | Great Cormorant | 9.7% |
| Australasian Grebe | 69.4% | Little Black Cormorant | 43.6% |
| Hoary-headed Grebe | 4.8% | Australasian Darter | 14.5% |
| White-headed Pigeon | 14.5% | Black-shouldered Kite | 1.6% |
| Brown Cuckoo-Dove | 21.0% | Square-tailed Kite | 4.8% |
| Crested Pigeon | 1.6% | Wedge-tailed Eagle | 3.2% |
| Bar-shouldered Dove | 64.5% | Little Eagle | 1.6% |
| Wompoo Fruit-Dove | 1.6% | Swamp Harrier | 8.1% |
| Pheasant Coucal | 37.1% | Grey Goshawk | 4.8% |
| Eastern Koel | 6.5% | Collared Sparrowhawk | 1.6% |
| Channel-billed Cuckoo | 1.6% | White-bellied Sea-Eagle | 22.6% |
| Horsfield's Bronze-Cuckoo | 4.8% | Whistling Kite | 30.7% |
| Shining Bronze-Cuckoo | 33.9% | Rainbow Bee-eater | 6.5% |
| Fan-tailed Cuckoo | 45.2% | Dollarbird | 9.7% |
| Brush Cuckoo | 17.7% | Azure Kingfisher | 24.2% |
| Pallid Cuckoo | 1.6% | Forest Kingfisher | 8.1% |
| Tawny Frogmouth | 1.6% | Sacred Kingfisher | 25.8% |
| White-throated Nightjar | 1.6% | Laughing Kookaburra | 66.1% |
| Australian Owlet-nightjar | 1.6% | Brown Falcon | 1.6% |
| White-throated Needletail | 8.1% | Yellow-tailed Black-Cockatoo | 9.7% |
| Purple Swamphen | 56.5% | Galah | 1.6% |
| Dusky Moorhen | 29.0% | Little Corella | 1.6% |
| Eurasian Coot | 67.7% | Australian King-Parrot | 6.5% |
| Black-winged Stilt | 1.6% | Red-rumped Parrot | 1.6% |
| Black-fronted Dotterel | 1.6% | Eastern Rosella | 59.7% |
| Masked Lapwing | 4.8% | Rainbow Lorikeet | 19.4% |
| Comb-crested Jacana | 74.2% | Scaly-breasted Lorikeet | 25.8% |
| Latham's Snipe | 11.3% | Regent Bowerbird | 6.5% |
| Black-necked Stork | 1.6% | Satin Bowerbird | 8.1% |

| | | | |
|-----------------------------|-------|----------------------------|-------|
| White-throated Treecreeper | 37.1% | Eastern Whipbird | 46.8% |
| Variegated Fairy-wren | 67.7% | Australasian Figbird | 11.3% |
| Superb Fairy-wren | 93.6% | Olive-backed Oriole | 41.9% |
| Red-backed Fairy-wren | 14.5% | Pied Currawong | 1.6% |
| Southern Emu-wren | 22.6% | Australian Magpie | 48.4% |
| Scarlet Honeyeater | 61.3% | Pied Butcherbird | 46.8% |
| Striped Honeyeater | 54.8% | Grey Butcherbird | 53.2% |
| Noisy Friarbird | 41.9% | White-breasted Woodswallow | 54.8% |
| Brown Honeyeater | 56.5% | Spangled Drongo | 12.9% |
| White-cheeked Honeyeater | 30.7% | Willie Wagtail | 77.4% |
| White-naped Honeyeater | 1.6% | Rufous Fantail | 9.7% |
| Eastern Spinebill | 64.5% | Grey Fantail | 85.5% |
| Lewin's Honeyeater | 88.7% | Torresian Crow | 22.6% |
| Little Wattlebird | 22.6% | Forest Raven | 19.4% |
| Yellow-faced Honeyeater | 61.3% | Australian Raven | 9.7% |
| Noisy Miner | 53.2% | Leaden Flycatcher | 8.1% |
| Spotted Pardalote | 14.5% | Restless Flycatcher | 4.8% |
| Striated Pardalote | 1.6% | Magpie-lark | 8.1% |
| Brown Gerygone | 83.9% | Black-faced Monarch | 12.9% |
| White-throated Gerygone | 25.8% | Eastern Yellow Robin | 85.5% |
| White-browed Scrubwren | 40.3% | Mistletoebird | 69.4% |
| Large-billed Scrubwren | 6.5% | Chestnut-breasted Mannikin | 4.8% |
| Yellow Thornbill | 67.7% | Red-browed Finch | 91.9% |
| Striated Thornbill | 1.6% | Golden-headed Cisticola | 1.6% |
| Brown Thornbill | 85.5% | Tawny Grassbird | 27.4% |
| Varied Sittella | 6.5% | Little Grassbird | 1.6% |
| Black-faced Cuckoo-shrike | 21.0% | Australian Reed-Warbler | 6.5% |
| White-bellied Cuckoo-shrike | 1.6% | Fairy Martin | 8.1% |
| Cicadabird | 16.1% | Tree Martin | 3.2% |
| Rufous Whistler | 41.9% | Welcome Swallow | 62.9% |
| Golden Whistler | 58.1% | Silvereye | 43.6% |
| Grey Shrike-thrush | 59.7% | Bassian Thrush | 1.6% |
| Crested Shrike-tit | 1.6% | | |

The 24 most commonly recorded species

| | | | |
|----------------------|-------|-------------------------|-------|
| Superb Fairy-wren | 93.6% | Australasian Grebe | 69.4% |
| Pacific Black Duck | 91.9% | Mistletoebird | 69.4% |
| Red-browed Finch | 91.9% | Eurasian Coot | 67.7% |
| Lewin's Honeyeater | 88.7% | Variegated Fairy-wren | 67.7% |
| Brown Thornbill | 85.5% | Yellow Thornbill | 67.7% |
| Grey Fantail | 85.5% | Laughing Kookaburra | 66.1% |
| Eastern Yellow Robin | 85.5% | Bar-shouldered Dove | 64.5% |
| Brown Gerygone | 83.9% | Eastern Spinebill | 64.5% |
| Black Swan | 80.7% | Welcome Swallow | 62.9% |
| Willie Wagtail | 77.4% | Chestnut Teal | 61.3% |
| Comb-crested Jacana | 74.2% | Scarlet Honeyeater | 61.3% |
| Grey Teal | 69.4% | Yellow-faced Honeyeater | 61.3% |